

ANALYZING THE IMPACT OF ACTIVE AND PASSIVE TOBACCO SMOKING ON RESPIRATORY HEALTH OF RURAL WOMEN RESIDING IN THE KANPUR REGION

ALKA KATIYAR¹, KAMINI JAIN², RAVISH KATIYAR³, NEELMA KUNVAR⁴,
NEENA SRIVASTAVA⁵ & POOJA GAUR³

¹Krishi Vigyan Kendra, Thariaon, Fatehpur, Uttar Pradesh, India

²Department of Food and Nutrition, Government PG Girls College, Itarsi, Madhya Pradesh, India

³Department of Microbiology, Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow, Uttar Pradesh, India

⁴Department of Extension, CSA University of Agriculture and Technology, Kanpur, Uttar Pradesh, India

⁵Department of Physiology, King George's Medical University, Lucknow, Uttar Pradesh, India

ABSTRACT

Inhaling the tobacco smoke is known health hazard that may cause many serious respiratory disorders such as COPD, asthma, lung cancer, bronchitis, etc., especially among rural women of low and middle-income countries. Continuous exposure of passive tobacco smoke may also adversely affect the respiratory health. In this study, total 145 rural women of 20-60 years age group, those involved in tobacco smoking, were selected from the Kanpur region. A high prevalence (N=121, 83.4%) of passive smoking was found among total tobacco smoking rural women. The highest percentage (54%) of active smokers was found between 50 to 60 years age group. Agricultural women showed the highest involvement in active tobacco smoking (42%) while the passive tobacco exposure was found highest among factory workers (28%). Bidi was found most popular tobacco smoking pattern among rural women (66%). Clinical manifestations such as coughing, sputum formation, difficulty in breathing, weakness, sore throat, headache, were found significantly higher among active smokers. The study concludes that the rate of passive smoking is very high among tobacco smoking rural women. Active smoking is most common among agricultural women while least common among students. Clinical manifestations were found significantly higher among active smokers than passive smokers.

KEYWORDS: Tobacco Smoking, Respiratory Health, Rural Women & Tobacco Patterns

Received: Oct 24, 2018; **Accepted:** Nov 14, 2018; **Published:** Nov 26, 2018; **Paper Id.:** IJEEFUSDEC20184

INTRODUCTION

Tobacco smoke is one of the most hazardous air pollutants that causes multiple health disorders and associated with large numbers of deaths worldwide (Rebase *et al.*, 1980, Nazaroff *et al.*, 2004) Approximately 80 % deaths associated with tobacco consumption occur in developing countries (Kulkarni *et al.*, 2017). In India, tobacco smoking is more common among rural women, about 0.5 % of women in urban areas and 2%, in rural areas used tobacco smoking. According to a previous study, female are comparatively more sensitive to tobacco smoke than male (Paoletti *et al.*, 1996).

Tobacco smoke contains more than 4,000 substances; some of them are cytotoxic and carcinogenic (Church *et al.*, 1985). Nicotine in tobacco smoke is highly addictive (Vijayakumar *et al.*, 2014). Tobacco smoking causes many health problems, but respiratory disorders such as - COPD, asthma, lung cancer, bronchitis etc. are more common among tobacco smokers. Over 75% of chronic bronchitis and 90% of lung cancer are associated with tobacco smoking (Murray *et al.*, 1997, Schrek *et al.*, 1950). Smoking is also found associated with other respiratory illness such as pneumonia, tuberculosis and flu-like illness (Bello *et al.*, 2014, Boon *et al.*, 2005, Wong *et al.*, 2013). In India, more than 60% of respiratory diseases are associated with tobacco smoking (Vijayakumar *et al.*, 2014). The effect of tobacco smoking on the respiratory health of rural women in the North Indian region is not well studied. The aim of this study is to analyze the impact of tobacco smoke on the respiratory health of rural women of Kanpur region.

MATERIALS AND METHODS

A cross-sectional study was conducted on 145 tobacco smoking rural women aged between 20-60 yrs, residing in rural areas of the Kanpur, Uttar Pradesh, India. The sample size determined using the formula $4pq/d^2$. Where the prevalence of tobacco smoking women in rural areas as 10%, relative error assumed to be 5%, a sample size was calculated 145. A pre-tested interview schedule was applied to collect the data. All women were asked questions from the prepared questionnaire about tobacco use, socioeconomic conditions and respiratory complications.

Inclusion Criteria

All women of 20-60 years age group with tobacco smoke exposure minimum 5 times a day continuously for the last three years.

Exclusion Criteria

Women those consumed tobacco occasionally or consumed smokeless tobacco were excluded from this study.

Clinical Manifestation

Common clinical manifestations of respiratory illness such as a cough, chest pain, shortness of breath, sputum formation, difficulty in breathing, sneezing, runny nose, weakness, headache, mild fever etc. were included in this study.

Statistical Analysis

Statistical analysis was done by using SPSS software (IBM SPSS statistics 20). The Chi-square test was performed to compare clinical manifestations of respiratory illness between active tobacco smokers and passive tobacco smokers.

RESULTS

From the study, it was found that there is a high prevalence (N=121, 83.4%) of passive smokers among a total 145 tobacco smoking rural women in Kanpur region. The highest percentage (54%) of active smokers was found between 50 to 60 years age group while the highest percentage (38%) of passive smokers belonged to 40-49 years age group (Figure 1). Agricultural women showed highest involvement in active tobacco smoking (42%) followed by factory workers (23%), housewives (19%), students showed the least involvement in active tobacco smoking while the passive tobacco exposure was found highest (28%) among factory workers (Figure 2). Inactive tobacco smoking patterns, bidi was found most popular tobacco smoking pattern among rural women (66%) followed by chilam (17%), hukka (13%) and cigarette (4%)

(Figure 3). Comparing the active smokers and passive smokers for clinical manifestations of respiratory illness such as coughing (67% vs 31%), sputum formation (47% vs 16%), difficulty in breathing (25% vs 9%), weakness (47% vs 21 %), sore throat (25% vs 9%), headache (29% vs 11%) were found significantly (p-value <0.05) higher among active smokers than passive smokers (Table 1).

Table 1: Clinical Manifestations Associated with Respiratory Illness among Tobacco Smoking Rural Women (n=145).

S.N.	Clinical Manifestations	Active Smokers (N=24)	Passive Smokers (N=121)	p Value*
1.	Coughing	16 (67.0%)	38 (31.0 %)	0.002
2.	Chest Pain	5 (21.0%)	16 (13.0 %)	0.346
3.	Difficulty in breathing	6 (25.0%)	11 (9.0 %)	0.038
4.	Shortness of breath	5 (21.0%)	9 (7.0 %)	0.057
5.	Sneezing	3 (13.0%)	10 (8.0 %)	0.452
6.	Runny nose	2 (8.0%)	7 (6.0%)	0.644
7.	Sputum formation	11 (47.0%)	19 (16.0 %)	0.001
8.	Weakness	11 (47.0%)	26 (21.0 %)	0.02
9.	Mild fever	4 (17.0%)	6 (5%)	0.061
10.	Headache	7 (29.0%)	13 (11%)	0.025
11.	Sore throat	6 (25.0%)	11 (9.0 %)	0.036

* Chi-square test was performed to compare clinical manifestation of respiratory illness between active tobacco smokers and passive tobacco smokers (p-value <0.05 was considered significant).




Figure 1: Age-Wise Percentage of Rural Women Involved in Active and Passive Tobacco Smoking




Figure 2: Effect of Education and Occupation of Rural Women on Tobacco Smoking




Figure 3: Percentage of Tobacco Smoking Patterns

DISCUSSIONS

The tobacco smoking by rural women has many consequences not only for their health but also for economic wellbeing, especially in low- middle-income countries. Women face various tobacco smoking associated health hazards like reproductive complications, multiple types of cancers and cardiovascular risks (Goel *et al.*, 2014). Most common adverse effect of tobacco smoking is found in the form of respiratory illness such as COPD, asthma, lung cancer, bronchitis etc. (Vijayakumar *et al.*, 2014, Schrek *et al.*, 1950). An increasing prevalence of tobacco smoking among rural women is a matter of serious public health concern. Women involved in active tobacco smoking not only affect their health but also cause a threat of passive smoking to other family members. In our study, a high prevalence (N=121, 83.4%) of passive smokers was found among total tobacco smoking, women (N=145). According to a previous study in China, the prevalence of passive smoking among rural areas found 63.4% (Li *et al.* 2015) while another study stated the exposure rate of passive smoking on the women of reproductive age was 43.46% (Gong *et al.*, 2016) High involvement of women between 50 to 60 years age group was found in this study, that is supported by a previous study performed in the same region (Katiyar *et al.*, 2017). In our study students showed the least involvement in both active and passive smoking. A previous Study supported our results showing least tobacco consumption among educated women. (Kathirvel *et al.*, 2014)

Inactive tobacco smoking patterns, bidi was found most popular tobacco smoking pattern among rural women (66%). According to a study, 84.1% of rural women of Bihar used bidi for smoking. (Sinha *et al.*, 2003) Clinical manifestations such as coughing, sputum formation, difficulty in breathing, weakness, sore throat, headache, were found significantly higher among active smokers than passive smokers. Similarly, according to a previous study, the active tobacco smoke exposure was significantly more prevalent among acute respiratory distress syndrome patients compared to passive smoking, (Hasieh *et al.*, 2014) while another study suggested that both active and passive smoking can cause respiratory illness. (Lam *et al.*, 1998).

CONCLUSIONS

The rate of passive smoking is very high among tobacco smoking rural women in Kanpur region. The involvement of older rural women in tobacco smoking is higher than women of the early age group. Active smoking is most common among agricultural women while least common among students. Clinical manifestations such as coughing, sputum formation, difficulty in breathing, weakness, sore throat, headache found significantly higher among active smokers than passive smokers.

ACKNOWLEDGEMENTS

We are thankful to Krishi Vigyan Kendra, Thariaon, Fatehpur (under C.S.A. University of Agriculture and Technology, Kanpur) for valuable support.

REFERENCES

1. Bello, S., Menéndez, R., Antoni, T., Reyes, S., Zalacain, R., Capelastegui, A. (2014). Tobacco smoking increases the risk for death from pneumococcal pneumonia. *Chest*, 146: 1029-1037.
2. Boon, S. D., Lill, S. W. P., Borgdorff, M. W., Verver, S., Bateman, E. D., Lombard, C. J., Enarson, D. A., Beyers, N. (2005). Association between smoking and tuberculosis infection: a population survey in a high tuberculosis incidence area. *Thorax*, 60: 555-557.
3. Church, D. F., Pryor, W. A. (1985). Free-radical chemistry of cigarette smoke and its toxicological implications. *Environ Health Perspect*, 64: 111-126.
4. Goel, S., Tripathy, J. P., Singh, R. J. and Lal, P. (2014). Smoking trends among women in India: Analysis of nationally representative surveys (1993-2009). *South Asian J Cancer*, 3: 200-202.
5. TAPRE, P., & Patel, H. R. (2015). Effect of bidi tobacco dust on root-knot nematode (*Meloidogyne incognita*) attacking okra. *Int. J. Agril. Sci. Res.* 5 (2): 55-58. s.
6. Gong, X., Luo, X. and Ling, L. (2016). Prevalence and Associated Factors of Secondhand Smoke Exposure among Internal Chinese Migrant Women of Reproductive Age: Evidence from China's Labor-Force Dynamic Survey. *Int J Environ Res Public Health*, 13: 371-381.
7. Hsieh, S. J., Zhuo, H., Benowitz, N. L., Thompson, B. T., Liu, K. D, Matthay, M. A., Calfee, C. S. (2014) Prevalence and Impact of Active and Passive Cigarette Smoking in Acute Respiratory Distress Syndrome. *Crit Care Med*, 42: 2058-2068.
8. Kathirvel, S., Thakur, J. S., Sharma, S. (2014). Women and tobacco: A cross sectional study from North India. *Indian Journal of Cancer*, 51: 78-82.

9. Katiyar, A., Jain, K., Kunvar, N., Katiyar, R., Gaur, P., Vaish, S. (2017). The study of oral cancer symptoms among tobacco consuming rural women in kanpur region. *World Journal of Pharmaceutical Research*, 6: 1268-1273.
10. Kulkarni, N.(2017). Women and Tobacco- An Overview. *Curr Trends Biomedical Eng & Biosci*, 2: 001-006.
11. Lam, T. H., Chung S. F., Betson C. L., Wong, C. M. and Hedley, A. J. (1998). Respiratory symptoms due to active and passive smoking in junior secondary school students in Hong Kong. *International Journal of Epidemiology*, 27: 41-48.
12. Li, Z., Yao, Y., Yu, Y., Shi, J., Liu, Y., Tao, Y. (2015). Prevalence and Associated Factors of Passive Smoking among Women in Jilin Province, China: A Cross-Sectional Study. *Int J Environ Res Public Health*, 12: 13970–13980.
13. Murray, C. J. L., Lopez, A. D. (1997). Alternative projections of mortality and disability by cause 1990–2020: global burden of disease. *Lancet*, 349: 1498–1504.
14. Nazaroff, W. W., Singer, B. C. (2004). Inhalation of hazardous air pollutants from environmental tobacco smoke in US residences. *J Expo Anal Environ Epidemiol*, 14: S71-S77.
15. Paoletti, P., Fornai, E., Maggiorelli, F., et al. (1996). Importance of base-line cotinine plasma values in smoking cessation: results from a double-blind study with nicotine patch. *Eur Respir J*, 9: 643-5.
16. Repace, J. L., Lowrey, A. H. (1980). Indoor air-pollution, tobacco-smoke, and publichealth. *Science*, 208: 464-472.
17. Schrek, R., Baker, L. A., Ballard, G. P., Dolgoff, P. (1950). Tobacco smoking as an etiologic factor in disease: cancer. *Cancer Res*, 10: 49–58.
18. Sinha, D. N., Gupta, P. C, Pednekar, M. S. (2003) Tobacco use in a rural area of Bihar, India. *Indian Journal of Community Medicine*, 28: 4.
19. Vijayakumar, A., Sreekanthan, K., Belicita, A. (2014). Respiratory Problems Among Smokers in a Rural Area in South India: A Pilot Study. *Indian Journal of Clinical Practice*, 25: 381-385.
20. Wong, C. M., Yang, L., Chan, K. P., Chan, W. M., Song, L., Lai, H. K., Thach, T.Q., Ho, L. M. (2013). Cigarette smoking as a risk factor for influenza associated mortality: evidence from an elderly cohort. *Influenza and Other Respiratory Viruses*, 7: 531–539.